

WHAT IS CLAIMED IS:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

(b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-976.

10 2. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 3 or SEQ ID NO: 6, optionally further comprising an amino-terminal methionine;

15 (b) an amino acid sequence for an ortholog of either SEQ ID NO: 2 or SEQ ID NO: 5;

(c) an amino acid sequence that is at least about 70 percent identical to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

20 (d) a fragment of the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic; and

25 (e) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-976, or the amino acid sequence of any of (a) - (c).

[illegible]

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

10 (c) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(d) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 that has a C- and/or N- terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

(e) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

4. An isolated polypeptide encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4;

(b) the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-976;

(c) a nucleotide sequence encoding the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

5

Sub
a1
comp

10

[illegible]

15

20

25

6. An isolated polypeptide encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(b) a nucleotide sequence encoding a polypeptide as set forth in either
5 SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion;

(d) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 that has a C- and/or N- terminal truncation;

(f) a nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides; and

20

25

8. A composition comprising the polypeptide of any of Claims 1, 2, or 3, and a pharmaceutically acceptable formulation agent.

9. The composition of Claim 8, wherein the pharmaceutically acceptable formulation agent is a carrier, adjuvant, solubilizer, stabilizer, or anti-oxidant.

Sub a2
5 10. ~~The composition of Claim 8, wherein the polypeptide comprises the amino acid sequence as set forth in either SEQ ID NO: 3 or SEQ ID NO: 6.~~

11. A polypeptide comprising a derivative of the polypeptide of any of Claims 1, 2, or 3.

10 12. The polypeptide of Claim 11 that is covalently modified with a water-soluble polymer.

Sub a3
15 13. ~~The polypeptide of Claim 12, wherein the water-soluble polymer is selected from the group consisting of polyethylene glycol, monomethoxy-polyethylene glycol, dextran, cellulose, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide copolymers, polyoxyethylated polyols, and polyvinyl alcohol.~~

20 14. A fusion polypeptide comprising the polypeptide of any of Claims 1, 2, or 3 fused to a heterologous amino acid sequence.

15. The fusion polypeptide of Claim 14, wherein the heterologous amino acid sequence is an IgG constant domain or fragment thereof.

Sub a4
25 16. A polypeptide produced by a process comprising culturing a host cell comprising a vector comprising a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4;

(b) the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-976;

(c) a nucleotide sequence encoding the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

5 (d) a nucleotide sequence that hybridizes under at least moderately stringent conditions to the complement of any of (a) - (c);
under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.

10 17. A polypeptide produced by a process comprising culturing a host cell comprising a vector comprising a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide that is at least about 70 percent identical to the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID
15 NO: 5, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4, the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-976, or the nucleotide sequence of (a);

(c) a region of the nucleotide sequence of either SEQ ID NO: 1 or SEQ ID NO: 4, the DNA insert in ATCC Deposit No. PTA-976, the nucleotide sequence (a) or (b), encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide fragment has an activity of the encoded polypeptide as set
20 forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic;

(d) a region of the nucleotide sequence of either SEQ ID NO: 1 or SEQ ID NO: 4, the DNA insert in ATCC Deposit No. PTA-976, or the nucleotide sequence of any of (a) - (c), comprising a fragment of at least about 16 nucleotides; and

See a 4 cont

(e) a nucleotide sequence that hybridizes under at least moderately stringent conditions to the complement of any of (a) - (d);
under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.

5

18. A polypeptide produced by a process comprising culturing a host cell comprising a vector comprising a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in either
10 SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(b) a nucleotide sequence encoding a polypeptide as set forth in either
15 SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(c) a nucleotide sequence encoding a polypeptide as set forth in either
20 SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(d) a nucleotide sequence encoding a polypeptide as set forth in either
SEQ ID NO: 2 or SEQ ID NO: 5 that has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

25 (e) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the encoded

10037960-0902000

Sub
a4
cont

an activity of
nucleotide se
nucleotides; and
nucleotide s
ons to the com
conditions to
n the culture.

5

under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.

15

20. The polypeptide of any of Claims 16, 17, or 18, wherein the host cell is a prokaryotic cell.

[illegible]